



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.


Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let P represent the stream or pond and BA ,² a portion of the line. With the instrument at the station S , set a flag at A , on the opposite side of the stream or pond, and also set a flag at D , any convenient point on the line, in the opposite direction from S . Take any point C , at right-angles with the line AB from the station S and at any convenient distance, which need not be measured. Set the instrument at the station C and direct the sights to the flag at A and note the angle SCA ; turn the instrument towards a flag to be placed at a point B , which must be kept in range with the flag at D by an assistant at S , making the angle SCB equal the angle SCA .



Now because of the similar triangles SCA and SCB having a common side CS , we have $SB = SA$; therefore chain back from S to B , and the result will be the length of the line produced to A .

When a back sight to B is not practicable the equivalent of the line SA may be chained by constructing on SA an equilateral triangle one side of which shall lie wholly on the opposite side of the stream.

QUERY. BY THE EDITOR. If a cube be cut by four vertical planes each of which intersects two contiguous vertical faces along their center lines, thus cutting away half the volume of the cube; and if the two ends of the remaining parallelopipedon be each cut by four planes, each of the four planes passing through the center of the upper and the lower base, respectively, making an angle of 45° with the vertical axis and intersecting two contiguous sides of the parallelopipedon at equal angles, thus cutting off the *eight corners* of the parallelopipedon, the remaining solid is a crystalline form known to mineralogists as a "rhombic dodecahedron," (Dana,) having twelve faces and fourteen solid angles. Through the center of this solid let a plane be passed which shall intersect its six parallel edges at right-angles, and the solid will be divided into two equal and similar solids which are identical in form with the honey bee's cell; that is, each will be a hexagon terminated by three planes which make respectively an angle of $54^\circ 44' 8''$ with the axis of the hexagon. Though this is the angle which gives a minimum surface for a given volume, and therefore would seem to indicate intelligence and design in its construction, yet being a crystalline form, may not the physical forces which determine the position of the molecules in the formation of the rhombic dodecahedron influence the action of the bee in the construction of its cell rather than the intelligence and design of the bee?